

Tracking the Progress 2022:
**National Sexually
Transmissible Infections
Strategy**



UNSW
SYDNEY



UNSW
Kirby Institute

The years for comparison in this report are from the end of 2017 to the end of 2021 unless focus is given to the impact of the COVID-19 epidemic, where the years for comparison are 2016 to 2019, and from 2019 to 2021.

Sources of data are provided in the data dashboard on the Kirby data site.

Acknowledgement is given to the many contributors helping report progress against the National Sexually Transmissible Infections Strategy. The full list of contributors can be found on the Acknowledgement page of the Kirby data site.

© Kirby Institute, UNSW Sydney 2023

ISSN 2653-2158 (Online)

This publication is available at Internet address
<http://www.kirby.unsw.edu.au>

Kirby Institute, UNSW Sydney

Suggested citation:

King, J., and S. McGregor. Tracking the Progress 2022: Executive Summaries. Kirby Institute, UNSW Sydney

Telephone: 02 9385 0900 Facsimile: 02 6100 2860 International prefix: 61 2
Email: info@kirby.unsw.edu.au

Tracking the Progress 2022: National Sexually Transmissible Infections Strategy

The goals of the Fourth National Sexually Transmissible Infections Strategy are to:

1. Reduce transmission of, and morbidity and mortality associated with STI in Australia
2. Eliminate the negative impact of stigma, discrimination and legal and human rights issues on people's health
3. Minimise the personal and social impact of STI

The National Sexually Transmissible Infections Strategy has five targets that provide specific focus for the efforts made towards achieving the goals of this Strategy. Each target has corresponding indicators that measure progress towards attaining the target. The full list of the targets and their indicators are listed in Appendix i.

The COVID-19 pandemic resulted in restrictions of access to healthcare, including testing and treatment. These restrictions likely impacted on progress against many indicators between 2019 and 2021. For this reason, the years for comparison in this report are from 2017 to 2021, and where appropriate, 2017 to 2019 and 2019 to 2021.

Target 1

access full
data here



Achieve and maintain HPV adolescent vaccination coverage of 80%

- HPV adolescent vaccine course completion is defined as the receipt of two vaccine doses if dose two is given at least five months after dose one. The definition of course completion is also met if three doses are given when dose two was given less than five months after dose one. The proportion of 15-year-old females registered in the Australian Immunisation Register receiving two HPV vaccine doses met the target in 2020 (81%) and 2021 (80%). In the same period, the proportion of 15-year-old males registered in the Australian Immunisation Register receiving at least two HPV vaccine doses increased from 70% in 2016 to 77% in 2019 then remained steady to be 77% in 2021.

Target 2

access full
data here



Reduce the prevalence of gonorrhoea, chlamydia and infectious syphilis

Part A: Notifications and testing

Gonorrhoea

- Between 2017 and 2019, the gonorrhoea notification rate increased by 20% from 118.3 to 141.4 Per 100 000. Between 2019 and 2021 the gonorrhoea notification rate declined by 23% from 141.4 to 109.4 per 100 000. Similar trends were seen among males and females and in 2021, notification rates were 149.1 and 69.4 per 100 000 population, respectively. Declines between 2019 and 2021 were likely attributable in part to the impact of COVID-19 restrictions on social activity, healthcare access and testing, and travel.
- Among people aged 15 to 29 years, the proportion of gonorrhoea tests yielding a positive test result between 2017 and 2021 fluctuated between 2.4% (in 2021) and 2.8% (in 2019). In this age group, similar trends were seen among males and females and in 2021, the proportion of gonorrhoea tests yielding a positive test result was 5.6% and 1.3%, respectively. The proportion of tests yielding a positive test result was higher among males than among females for every year reported.
- Among female sex workers attending sexual health clinics participating in the Australian Collaboration for Coordinated Enhanced Sentinel Surveillance (ACCESS), the incidence of gonorrhoea fluctuated between 2017 and 2021 and was 7.6 per 100 person years in 2021.
- Between 2017 and 2021, among gay and bisexual men attending clinics participating in ACCESS, the incidence of gonorrhoea fluctuated between 2017 and 2019 but declined by 18% between 2019 and 2021 from 26.8 to 22.0 infections per 100 person-years.

Target 2

access full
data here



Reduce the prevalence of gonorrhoea, chlamydia and infectious syphilis

Part A: Notifications and testing

Chlamydia

- Between 2017 and 2019, the chlamydia notification rate remained stable. Between 2019 and 2021 the chlamydia notification rate declined by 17% from 435.1 to 362.7 per 100 000. Similar trends were seen among males and females and in 2021 notification rates were 344.0 and 383.7 per 100 000, respectively. Declines in 2020 are likely attributable in part to the impact of COVID-19 restrictions on social activity, healthcare access and testing, and travel.
- Between 2017 and 2021, among people aged 15 to 29 years, the proportion of chlamydia tests yielding a positive test result fluctuated between 13.0% (in 2017 and 2018) and 11.8% (in 2021).
- Among female sex workers attending clinics participating in ACCESS, the chlamydia incidence fluctuated between 11.4 and 11.0 new infections per 100 person-years and was 11.0 new infections per 100 person-years in 2021.
- Among gay and bisexual men attending clinics participating in ACCESS, the incidence of chlamydia increased by 15% between 2017 and 2021 from 26.4 to 30.3 new infections per 100 person-years.

Infectious syphilis

- Between 2017 and 2019 the infectious syphilis notification rate increased by 30% from 18.4 to 23.9 notifications per 100 000 and then fluctuated between 2019 and 2021. In 2021, the infectious syphilis notification rate was 22.7 per 100 000. In the same period, similar trends were seen among males and females and in 2021, notification rates were 37.0 and 8.5 per 100 000, respectively. The changing in trends from 2020 were likely attributable in part to the impact of COVID-19 restrictions on social activity, healthcare access and testing, and travel.
- Between 2017 and 2021, among female sex workers attending sexual health clinics participating in ACCESS, the incidence of infectious syphilis fluctuated between 0.2 (in 2018 and 2021) and 0.5 (in 2020) new infections per 100 person-years.
- Among gay and bisexual men attending clinics participating in ACCESS, the incidence of infectious syphilis increased between 2017 and 2021 by 31% from 4.5 to 5.9 per 100 000 person-years.

Target 2

access full
data here



Reduce the prevalence of gonorrhoea, chlamydia and infectious syphilis

Part B: Knowledge and risk behaviours

- Among participants of the National Survey of Australian Secondary Students and Sexual Health (SASSH):
 - In 2018, 94.5% of students knew that someone could have a sexually transmissible infection without having any obvious symptoms, up from 89.4% of participants in 2013. This question was not asked in the 2021 survey.
 - In 2018, 72.8% of participants were aware that chlamydia affects both men and women, up from 60.7% of students in 2013. This question was adapted in 2021 to ask students if chlamydia is an STI that only affects persons with a womb/uterus. In 2021, 58.5% of respondents answered this question correctly.
 - In 2018, 53.6% of participants were aware that chlamydia can lead to infertility amongst women, down from 57.4% in 2013. In 2021, this question was adapted to ask if chlamydia can lead to infertility without specifying a population affected with 42% of participants answering this question correctly.
 - In 2021, 34.0% of participants knew that once a person caught genital herpes, they will always have the virus, down from 46.7% in 2013.
 - In all years and for all questions, a higher proportion of female students answered STI knowledge questions correctly than their male peers.
 - Overall, the highest levels of knowledge regarding STIs were demonstrated about the potentially asymptomatic nature of many infections, and lower levels of knowledge were seen in relation to chlamydia and herpes. Further detail on responses by gender is available on the [Kirby data site](#).
- The proportion of all sexually active respondents to the 2021 SASSH survey reporting always using a condom when they had sex in the last 12 months was 38.3%, a decrease from 46.3% in 2013. The proportion reporting condom use at last sex was 48.6% in 2021, a decline from 58.8% in 2013. Reported condom use was higher among males than females in all years.
- In 2018, almost a fifth (19.1%) of participants reported three or more sexual partners in the past year, a decrease from 26.4% in 2013.
- In 2021, the number of sexual partners was stratified by anal and vaginal sex.

Target 2

access full
data here



Reduce the prevalence of gonorrhoea, chlamydia and infectious syphilis

Part B: Knowledge and risk behaviours

- With 31.2% reporting three or more vaginal sexual partners (33.3% among females, 24.5% among males). Of those reporting anal sex, 9.2% reported three or more anal sex partners (3.9% among females, 19.7% among males).
- Full breakdowns of reported risky behaviours by sex are available on the [Kirby data site](#).
- Among participants of the Gay Community Periodic Survey (GCPS), the proportion who reported consistent condom use with casual sexual partners in the previous 12 months decreased steadily from 30.3% in 2017 to 17.0% in 2021.

Target 3

access full
data here



Increase STI testing coverage in priority populations

- Chlamydia and gonorrhoea are tested for concurrently in Australia using duplex testing. The proportion of 15- to 29-year-olds consulting with a general practitioner and receiving at least one chlamydia/gonorrhoea test from their general practitioner in the previous 12 months remained steady between 2016 and 2019 and was 14.3% in 2019. Between 2019 and 2021, this proportion increased from 14.3% to 16.7%. Among the same age group, similar trends were seen among both males and females, and in 2021, the proportion receiving at least one chlamydia/gonorrhoea test from their general practitioner in the previous 12 months was 9.6% and 22.6%, respectively. Despite the higher proportion of young people getting tested for chlamydia at their general practice, the overall lower number of tests conducted from 2020 suggests that the number of people seeking health care at their general practice declined from 2020.
- Between 2017 and 2021, among gay and bisexual men attending clinics participating in ACCESS, the proportion receiving a chlamydia, gonorrhoea and syphilis test in the previous 12 months declined steadily from 83.9% to 79.3%
- In the same period, among sex workers attending participating sexual health clinics in ACCESS, the proportion receiving a chlamydia, gonorrhoea and syphilis test in the previous 12 months increased from 87.4% in 2016 to 90.8% in 2019 and then declined to 80.2% in 2020.
- Among participants in the GCPS, the proportion who reported having had comprehensive STI testing in the previous 12 months increased steadily from 45.5% in 2017 to 57.1% in 2019, then declined to 47.1% in 2021.

Target 4

*access full
data here*



Eliminate congenital syphilis

- Between 2017 and 2021 the number of congenital syphilis notifications almost doubled, increasing from eight to 15 notifications. Similarly, when translated to notification rate, in the same period, the congenital syphilis notification rate almost doubled from 2.6 to 5.1 notifications per 100 000 live births.
- Among women of reproductive age (15 to 44 years), the infectious syphilis notification rate increased between 2016 and 2021 by 60% from 11.2 notifications to 17.9 notifications per 100 000.
- The large increase in the number of congenital syphilis cases between 2019 and 2021, despite little change in the notification rate, may mean there are large numbers of undiagnosed cases of infectious syphilis among women of reproductive age.

Target 5

access full
data here



Minimise the reported experience and expression of stigma in relation to STI

- Among the participants of the 2018 National Debrief Survey aged between 15 to 29 years:
 - 63% reported that they had never behaved negatively towards people with an STI while 8% reported that had sometimes or often behaved negatively towards someone with an STI. 28% reported they had done so rarely.
 - 12% reported that they would expect to never experience stigma if they had an STI while 62% reported they would sometimes, often or always expect to experience stigma. 26% reported they would expect to rarely experience stigma if they had an STI.
 - Among respondents with a history of having an STI, 50% reported that they had never experienced stigma or discrimination due to their STI while 27% reported they had sometimes, often or always experienced stigma or discrimination. 23% reported they had rarely experienced stigma or discrimination due to their STI.
- As reported in the ***Annual Report of Trends in Behaviour 2019 HIV and STIs in Australia***:
 - Among surveyed health care workers, 57% reported never witnessing negative behaviour towards people because of an STI. Conversely, 24% of health care workers reported they had sometimes, often, or always witnessed negative behaviour towards people because of an STI while 19% reported they had witnessed such behaviour rarely.
- As reported in the ***Annual Report of Trends in Behaviour 2022 HIV and STIs in Australia***:
 - Among surveyed health care workers in 2021, 57% reported never expressing negative behaviour towards people because of an STI down from 75% in 2018. Conversely, 22% of health care workers reported they had sometimes, often, or always expressed negative behaviour towards people because of an STI (up from 6% in 2018), while 19% reported that they had done so rarely (up from 18% in 2018).

Appendix

Sexually Transmissible Infections Strategy

	Indicator	Data source
Target 1	Achieve and maintain human papillomavirus adolescent vaccination coverage of 80%.	
	1a HPV two dose vaccination coverage for males and females aged 15 years of age.	Australian Immunisation Register (AIR)
Target 2	Reduce the prevalence of gonorrhoea, chlamydia and infectious syphilis.	
	2a Annual rate of gonorrhoea notifications.	National Notifiable Diseases Surveillance System (NNDSS)
	2b Annual rate of chlamydia notifications.	National Notifiable Diseases Surveillance System (NNDSS)
	2c Annual rate of infectious syphilis notifications.	National Notifiable Diseases Surveillance System (NNDSS)
	2d Incidence of STIs in sex workers attending a sexual health clinic.	ACCESS (Australian Collaboration for Coordinated Enhanced Sentinel Surveillance)
	2e Incidence of STIs in gay and bisexual men attending a health service.	ACCESS (Australian Collaboration for Coordinated Enhanced Sentinel Surveillance)
	2f Proportion of chlamydia tests that yield a positive result in the 15-29 year age group.	Medicare, National Notifiable Diseases Surveillance System (NNDSS)
	2g Proportion of gonorrhoea tests that yield a positive result in the 15-29 year age group.	Medicare, National Notifiable Diseases Surveillance System (NNDSS)
	2h Proportion of secondary school students giving the correct answer to STI knowledge and behaviour questions.	National Survey of Australian Secondary Students and Sexual Health
	2i Proportion of secondary school students reporting certain risky sexual behaviours.	National Survey of Australian Secondary Students and Sexual Health
	2j Proportion of young people (15-29 year olds) giving the correct answer to STI knowledge questions.	The Debrief Survey
	2k Proportion of young people (15-29 year olds) reporting consistent condom use with sexual partners in the previous 12 months.	The Debrief Survey
	2l Proportion of gay and bisexual men who reported consistent condom use with casual sexual partners in the previous 12 months.	Gay Community Periodic Surveys

Appendix

Sexually Transmissible Infections Strategy

	Indicator	Data source
Target 3	Increase STI testing coverage in priority populations.	
3a	Proportion of 15-29 year olds receiving at least one chlamydia test in the previous 12 months.	Medicare
3b	Proportion of 15-29 year olds receiving at least one gonorrhoea test in the previous 12 months.	Medicare
3c	Proportion of gay and bisexual men who report having had an STI test in the previous 12 months.	ACCESS (Australian Collaboration for Coordinated Enhanced Sentinel Surveillance)
3d	Proportion of gay and bisexual men attending a health clinic receiving a chlamydia, gonorrhoea and infectious syphilis test at least once in the previous 12 months.	ACCESS (Australian Collaboration for Coordinated Enhanced Sentinel Surveillance)
3e	Proportion of gay men who report having had comprehensive STI testing in the previous 12 months.	Gay Community Periodic Surveys
3f	Proportion of sex workers attending a health clinic receiving a chlamydia, gonorrhoea or infectious syphilis test in the previous 12 months	ACCESS (Australian Collaboration for Coordinated Enhanced Sentinel Surveillance)
3g	Proportion of young people (15-29 years) who reported having sex and have had an STI and/or HIV test in the previous 12 months.	The Debrief Survey
Target 4	Eliminate congenital syphilis.	
4a	Number of congenital syphilis notifications.	National Notifiable Diseases Surveillance System (NNDSS)
4b	Notification rate of congenital syphilis per 100 000 live births.	National Notifiable Diseases Surveillance System (NNDSS)
4c	Annual notification rate of infectious syphilis in women of reproductive age (15-44 years of age).	National Notifiable Diseases Surveillance System (NNDSS)
Target 5	Minimise the reported experience and expression of stigma in relation to STI.	
5a	Proportion of young people reporting negative behaviour towards people with an STI.	The Debrief Survey
5b	Proportion of people who report that they would expect to experience stigma if they had an STI.	The Debrief Survey
5c	Proportion of young people who report that they experienced stigma or discrimination due to their STI.	Annual Report of Trends in Behaviour
5d	Proportion of health care workers reporting or witnessing negative behaviour towards people with an STI.	Annual Report of Trends in Behaviour